

## To those who regularly take blood thinners

We are asking you the type of your medications in the pre-vaccination screening questionnaire because bleeding is difficult to stop if you are taking blood thinners.

You may receive the vaccine, but you will have to be cautious about any bleeding after the injection.

### Precautions for vaccination (intramuscular injection)



Please press the injected area firmly for at least 2 minutes.

- If you have symptoms like swelling in the arm, numbness in the arm, etc., please consult the doctor.
- You do not have to discontinue your medication before vaccination.

Medications  
that need  
precautions

### Anticoagulants

These are medications mainly prescribed to patients who have arrhythmia, thrombosis, or had surgery of the heart.

Product name	General Name
Warfarin	Warfarin Potassium
Prazaxa	Dabigatran Etxilate
Xarelto	Rivaroxaban
Eliquis	Apixaban
Lixiana	Edoxaban Tosilate Hydrate

There are other blood thinners such as below, but you may be vaccinated as usual.

### Antiplatelets, etc.

These are prescribed mainly for patients with arteriosclerosis, angina pectoris/myocardial infarction, after a cerebral infarction, or with arterial occlusion of the lower limb.

Product name	General Name	Product name	General Name
Bayaspirin	Aspirin	ComPlavin saCA	Clopidogrel Sulfate, Aspirin
Panaldine	Ticlopidine Hydrochloride	TAKELDA Combination Tablets	Aspirin Lansoprazole
Plavix	Clopidogrel Sulfate	Epadel	Ethyl Icosapentate
Effient	Prasugrel Hydrochloride	Dorner, Procylin	Beraprost Sodium
Pletal	Cilostazol	Anplag	Sarpogrelate Hydrochloride
Bufferin	Aspirin and Dealuminate composition		
Brilinta	Ticagrelor		etc.